

HARMFUL • ECOTOXIC

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

FIESTAR

NON-SELECTIVE HERBICIDE

ACTIVE INGREDIENT: 200 g/L GLUFOSINATE-AMMONIUM
in the form of a soluble concentrate

For non-residual control of a wide range of broadleaf and grass weeds
in orchards, vineyards, forestry and non-cropland areas.



Registered pursuant to ACVM Act 1997, Approval No. P8141
See www.foodsafety.govt.nz for registration conditions
Approved pursuant to HSNO Act 1996, Approval No. HSR000152
See www.epa.govt.nz for approval conditions

Kenso Corporation (M) Sdn Bhd Office A, 49B, Apollo Drive, Rosedale, Auckland 0632 New Zealand Ph: 09 410 0861

FIESTAR NON-SELECTIVE HERBICIDE

This product **MUST** be under personal control of an Approved Handler when applied in a wide dispersive manner, or when applied by a commercial contractor.

HSNO classification: 3.1D, 6.1D, 6.3B, 6.4A, 6.8B, 6.9A, 9.1D, 9.2A, 9.3C
HARMFUL

May be harmful if swallowed, inhaled or absorbed through the skin. May cause eye and mild skin irritation. Avoid contact with eyes and skin. Avoid inhalation of spray mist. May cause organ damage from repeated oral exposure at high doses.

ECOTOXICITY

Toxic to aquatic organisms. Avoid contamination of any water supply with product or empty container. Very toxic to soil flora.

BEWARE: APPLY THIS PRODUCT CAREFULLY. SPRAY DRIFT MAY CAUSE SERIOUS DAMAGE TO OTHER DESIRABLE PLANTS.

PRECAUTIONS

Do not eat, drink or smoke while using. Wear protective clothing such as impervious gloves, waterproof hat, coat and trousers when using. Remove protective clothing and wash hands and face thoroughly before meals and after work. May be harmful if swallowed, inhaled or absorbed through the skin. Avoid skin contact or inhalation. Avoid contact with the eyes.

STORAGE

Store in original container tightly closed and in a locked dry, cool, well ventilated area, away from foodstuffs. When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

FIRST AID

If swallowed do NOT induce vomiting. Take patient to hospital without delay. If skin contact occurs, remove contaminated clothing and flush skin with

running water, if irritation occurs seek medical advice. If concentrate is splashed in eye, flush with running water for at least 15 minutes. Call the National Poisons Centre 0800 POISON (0800 764 766) (24 hrs) or a doctor immediately for advice.

SAFETY DATA SHEET

A Safety Data Sheet is available from Kenso Corporation (M) Sdn Bhd on request.

CONDITIONS OF SALE

"Kenso Corporation (M) Sdn. Bhd." ("Kenso") shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Kenso's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Kenso has any authority to add to or alter these conditions.



EMERGENCY RESPONSE
Ring 0800 CHEMCALL, or
0800 243 622 (24 hours)

NET CONTENTS:



Product of Malaysia



Batch No.:

Date of Manufacture:

HARMFUL • ECOTOXIC

KEEP OUT OF REACH OF CHILDREN
READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

FIESTAR

NON-SELECTIVE HERBICIDE

**ACTIVE INGREDIENT: 200 g/L GLUFOSINATE-AMMONIUM
in the form of a soluble concentrate.**

For non-residual control of a wide range of broadleaf
and grass weeds in orchards, vineyards,
forestry and non-cropland areas.

Registered pursuant of ACVM Act 1997, Approval No. P8141
See www.foodsafety.govt.nz for registration conditions

Approved pursuant to HSNO Act 1996, Approval No.
HSR000152
See www.epa.govt.nz for approval conditions



Kenso Corporation (M) Sdn Bhd
Office A, 49B, Apollo Drive
Rosedale, Auckland 0632 NZ
Phone: (09) 410 0861

STORAGE

Store in original container tightly closed and in a
locked dry, cool, well ventilated area, away from food-
stuffs. When stored appropriately this product should
show no significant degradation for 2 years from the
date of manufacture. Contact your supplier for further
information about the use of any product that is older
than this.

SPILLS AND DISPOSAL

Contain the spill by damming, recover spilt product
by absorbing with sawdust or an inert absorbent
material then transfer the recovered spilt material to a
properly labeled drum. Deal with all spillages immedi-
ately. If contamination of drains, streams, water-
courses, etc. is unavoidable, warn the local water
authority. Dispose of this product only by using in
accordance with this label, or at an approved landfill,
or other approved facility. Do not burn. Triple rinse
container and add rinsate to the spray tank container
for any other purpose.

FIRST AID

If swallowed do NOT induce vomiting. Take patient to
hospital without delay. If skin contact occurs, remove

contaminated clothing and flush skin with running
water, if irritation occurs seek medical advice. If
concentrate is splashed in eye, flush with running
water for at least 15 minutes. Call the National
Poisons Centre 0800 POISON (0800 764 766) (24
hrs) or a doctor immediately for advice.

SAFETY DATA SHEET

A Safety Data Sheet is available from Kenso Corpora-
tion (M) Sdn Bhd on request.

CONDITION OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any
responsibility whatsoever and howsoever arising and
whether for consequential loss or otherwise in
connection with the supply of these goods other than
responsibility for the merchantable quality of the
goods and such responsibilities mandatorily imposed
by Statutes applicable to the sale or supply of these
goods. To the extent allowed by such Statutes the
liability of Kenso Corporation (M) Sdn. Bhd. is limited
to the replacement of the goods or (at the option of
Kenso Corporation (M) Sdn. Bhd.) the refund of the
price paid and where possible sufficient part of the
goods to enable proper examination being returned

DIRECTIONS FOR USE

Mixing:

Add the required amount of Fiestar to a partly filled spray tank with the agitation system operating. Complete filling the tank with water and apply, maintaining agitation.

Compatibility:

Fiestar is compatible with most commonly used residual herbicides.

WITHHOLDING PERIOD

28 days all crops

WEED SUSCEPTIBILITY

The lower rates indicated are appropriate when weeds are young, succulent and actively growing, or when the weed population is sparse and good coverage and penetration is possible. Use the higher rates when weeds are mature or the population is dense.

NB. Fiestar will not completely control mature weeds with well developed tap roots or underground storage organs: some re-growth may occur following treatment.

INDICATED RATE RANGE FOR VARIOUS WEED SPECIES

3-5 litres/ha in 300 - 500 litres of water:

Annual Poa, Asparagus seedlings, Beaked hawksbeard, Black nightshade, Calandrinia, Chickweed, Cleavers, Clover seedlings, Cornbind, Daisy, Dandelion, Mouse-ear, chickweed, Musky storksbill, Redroot, Rough-stalked, meadow grass, Scotch thistle, Sow thistle, Staggerweed, Summer grass, Spurrey, Stinking mayweed, Subterra nean clover, Wild turnip, Willow weed, Wireweed.

to Kenso Corporation (M) Sdn. Bhd. within thirty days
of delivery.

* Not a trademark of Kenso Corporation (M) Sdn Bhd.

Registered pursuant to ACVM Act 1997, Approval
No. P8141

See www.foodsafety.govt.nz for registration condons

Approved pursuant to HSNO Act 1996, Approval No.
HSR000152

See www.epa.govt.nz for approval conditions

Batch No. :

Date of Manufacture :

EMERGENCY RESPONSE
Ring 0800 CHEMCALL, or
0800 243 622 (24 hours)

This product MUST be under personal control of an Approved Handler when applied in a wide dispersive manner, or when applied by a commercial contractor.



HSNO classification: 3.1D, 6.1D, 6.3B, 6.4A, 6.8B, 6.9A, 9.1D, 9.2A, 9.3C

HARMFUL

May be harmful if swallowed, inhaled or absorbed through the skin. May cause eye and mild skin irritation. Avoid contact with eyes and skin. Avoid inhalation of spray mist. May cause organ damage from repeated oral exposure at high doses.

ECOTOXICITY

Toxic to aquatic organisms. Avoid contamination of any water supply with product or empty container. Very toxic to soil flora.

BEWARE: APPLY THIS PRODUCT CAREFULLY. SPRAY DRIFT MAY CAUSE SERIOUS DAMAGE TO OTHER DESIRABLE PLANTS.

PRECAUTIONS

Do not eat, drink or smoke while using. Wear protective clothing such as impervious gloves, waterproof hat, coat and trousers when using. Remove protective clothing and wash hands and face thoroughly before meals and after work. May be harmful if swallowed, inhaled or absorbed through the skin. Avoid skin contact or inhalation. Avoid contact with the eyes.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

5 - 7.5 litres/ha in 300 – 500 litres of water:

Bristle grass, Bur medick, Catsear, Chewing’s fescue, Clovers, Cut-leaved geranium, Docks, Fleabane, Parsley piert, Prairie grass, Ryegrasses, Sheep’s sorrel, Storksbill, Twin cress, Yorkshire fog.

7.5 – 10 litres/ha in 300 – 500 litres of water:

Broom, Buttercups Couch, Field pansy, Crowfoot grass, Gorse, Hawksbeard, Kikuyu grass, Mallows, Paspalum, Tall fescue, Wild strawberry, Willow herb, Yarrow.

10 litres plus/ha in 300 – 500 litres of water:

Bracken, Meadow rice grass, Plantains, Rape, Rushes, Sedges, Indian doab.

Resistant Weeds:

Hairy buttercup, some Ryegrass strains.

ADDITIONAL INFORMATION

Fiestar is absorbed into the foliage and green stems of actively growing plants. It is inactive in soil and is not translocated as an active herbicide, so thorough coverage of target weeds is essential for good results. Visual effects of treatment will become evident more rapidly under ideal growing conditions, generally within 3-10 days.

Rainfall within 2 hours of application may wash Fiestar off the foliage, and within 6 hours will reduce effectiveness to the degree that re-treatment may be necessary.

NB: Do not spray in temperatures below 10 oC. Avoid contact on green parts (including immature bark) of desirable plants, as damage may occur. Do not mow or cultivate after treatment until weeds start to die off.

RECOMMENDATIONS FOR USE:

Use	Rate	Remarks
Berryfruit-Rasperies and Brambles Primocane control	500-750ml/100 litres applied through cone nozzles in sufficient water to ensure complete coverage of primocanes.	Apply as a directed high volume spray to primocanes not exceeding 25cm in length. Avoid drift onto foliage of fruiting canes. Use the high rate on vigorously growing crops. The addition of Citowett* at 25ml/100 litres is recommended. “Knockdown” takes approximately 10 days. Do not apply to plants before the first primocanes for the season have reached 15cm high in height. Do not apply to plants under stress.
Orchard and Vineyard weed control. Pipfruit, Kiwifruit, Stonefruit, Grapes and Citrus.	Use a rate appropriate to the weeds present. If dilution rate is less than 1 litre/100 litres of water and Citowett* at 25ml/100 litres.	Apply in spring when the majority of the annual weeds have germinated. Residual herbicides may be applied in mixture for long-term weed control. Avoid contact with the foliage and immature bark of desirable plants.
Stale seedbed preparation particularly where clover is a problem weed.	3-5 litres/ha in 200-300 litres of water.	Apply to seedling weeds and clover in sufficient water to ensure complete coverage. May be applied up to one day before crop emergence.
Non-crop land. General weed control.	7.5-10 litres/ha in 300-500 litres of water, or 100ml/10 litre knapsack.	Apply in a drenching spray to the point of run-off. May be mixed with residual herbicides for long-term weed control.