

**TOXIC · ECOTOXIC · CORROSIVE**

**KEEP OUT OF REACH OF CHILDREN**

**READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING**

# **KEN-AMINE 625**

## **SELECTIVE HERBICIDE**

**ACTIVE INGREDIENT: 625 g/L 2,4-D as the dimethlyamine and the diethanolamine salts  
in the form of a soluble concentrate**

Non-volatile 2,4-D liquid for broadleaf weed control in cereals and pasture.



Registered pursuant of ACVM Act 1997, Approval No. P8709  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions  
Approved pursuant to HSNO Act 1996, Approval No. HSR000367  
See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions

# KEN-AMINE 625 SELECTIVE HERBICIDE

## APPROVED HANDLER

This product must be under the personal control of an approved handler at all times unless secured.

**HSNO CLASSIFICATION: 6.1C, 6.9A, 8.3A, 9.1B, 9.2A, 9.3B**

## TOXICITY

Toxic – may be fatal if swallowed, inhaled or absorbed through the skin. May cause organ damage from repeated oral exposure at high doses.

## CORROSIVE

Danger – this product is corrosive and may cause eye damage.

## ECOTOXIC

Toxic to aquatic organisms. Avoid contamination of any water supply with product or empty container.

Selective herbicide – very toxic to some plant species.

**Beware: Spray drift hazard. Apply this product carefully. Spray drift may cause serious damage to other desirable plants.** Ensure compliance with relevant regional air management plans. To minimise the risk of damage from drift of this product, especially where susceptible vegetation is within 0.5km, use low drift nozzles which minimise the production of small droplets (less than 100 microns). Maintain a water rate of at least 30 litres/ha for aerial spraying. Do not add additional wetter unless specified on the label. Preferred wind conditions for spraying are

a light constant breeze away from any sensitive crop.

Toxic to terrestrial vertebrates.

## FIRST AID

If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If splashed in eyes, wash out immediately with water. If inhaled move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped.

## SAFETY DATA SHEET

A Safety Data Sheet is available from Kenso Corporation (M) Sdn Bhd on request.

## STORAGE

Store in original container tightly closed and in a locked, dry, cool area away from foodstuffs, seeds, and fertilisers. Storage must be in accordance with NZS 8409 Management of Agrichemicals.

Stores containing 100 litres of Ken-Amine 625 Selective Herbicide require signage and secondary containment, and more than 100 litres require emergency response plans.

## CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.



**NET CONTENTS: 20L**



**EMERGENCY RESPONSE**  
Ring 0800 CHEMCALL, or  
0800 243 622 (24 hours)

Product of Malaysia



UN NO: 3082

ENVIRONMENTALLY HAZARDOUS  
SUBSTANCE, LIQUID N.O.S.

(CONTAINS 2,4-D 62.5%)

PACKING GROUP III

HAZCHEM 2X



In a transport emergency dial 111,  
Police or Fire Service.

For specialist advice in emergency only,  
call 0800 243 622 (24 hours)

**TOXIC • ECOTOXIC • CORROSIVE**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING**

# KEN-AMINE 625

## SELECTIVE HERBICIDE

**ACTIVE INGREDIENT: 625 g/L 2,4-D as the dimethylamine and the diethanolamine in the form of a soluble concentrate.**

Non-volatile 2,4-D liquid for broadleaf weed control in cereals and pasture.

Registered pursuant to ACVM Act 1997, Approval No. P8709  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions

Approved pursuant to HSNO Act 1996, Approval No. HSR000367  
See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions



**Kenso Corporation (M) Sdn Bhd**  
Office A, 49B, Apollo Drive  
Rosedale, Auckland 0632 NZ  
Phone: (09) 410 0861

### EQUIPMENT

Apply using well maintained spray equipment that is accurately calibrated to apply 100-300 litres of water/ha for ground spraying and 30-100 litres of water/ha for aerial spraying. Use higher water rates for larger weeds.

### DISPOSAL

Triple rinse empty container and add residue to spray tank. Recycle empty container through Agrecovery (0800 247 326, [www.agrecovery.co.nz](http://www.agrecovery.co.nz)). Otherwise, crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.

### Spillage:

Wear appropriate protective clothing and prevent material from entering waterways. Absorb spill with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely in an approved landfill.

### Transport:

Do not carry this product on a passenger service vehicle.

### FIRST AID

If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If splashed in eyes, wash out immediately with water. If inhaled move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped.

### SAFETY DATA SHEET

A Safety Data Sheet is available from Kenso Corporation (M) Sdn Bhd on request.

### CONDITION OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

\* Not a trademark of Kenso Corporation (M) Sdn Bhd.

Registered pursuant to ACVM Act 1997, Approval No. P8709  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions

Approved pursuant to HSNO Act 1996, Approval No. HSR000367  
See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions

Batch No. :

Date of Manufacture :

Stores containing 100 litres of Ken-Amine 625 Selective Herbicide require signage and secondary containment, and more than 100 litres require emergency response plans.

### PERSONAL PROTECTION

Avoid contact with skin or eyes and inhalation of spray mist.

When handling wear eye protection, waterproof gloves, boots and overalls. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work. Wash protective clothing daily after work.

### Tracking/Record Keeping:

This product requires tracking. The location and movement of this product must be recorded at each stage of its lifecycle as described in NZS 8409 Management of Agrichemicals.

**Shelf Life:** Shelf Life: When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

### Mixing:

Pour the measured quantity of Ken-Amine 625 Selective Herbicide into the partially filled spray tank and with the agitator running top up with the remainder of the water. Keep agitated during use.

### Cleaning:

Flush equipment thoroughly with clean water after use. Decontaminate before using on plants that are sensitive to 2,4-D.

### Decontamination:

After thoroughly washing the tank and flushing the system with clean water, quarter fill the tank, add 500g of an alkali detergent and circulate throughout the system for 15 minutes. Drain the system. Remove the filters and nozzles and clean them separately. Then flush the system with clean water and allow to drain.

**Compatibility:** Ken-Amine 625 Selective Herbicide is compatible with most other agricultural chemicals, including Ken-Met 600, Clopyralid 300g/L, Flumetsulam 800g/kg, Dicamba 500g/L and Bonza\*.

**EMERGENCY RESPONSE**  
Ring 0800 CHEMCALL, or  
0800 243 622 (24 hours)



**UN NO : 3082**  
**ENVIRONMENTALLY HAZARDOUS SUSTANCE,**  
**LIQUID N.O.S.**  
**(CONTAINS 2,4-D 62.5%)**  
**PACKAGING GROUP III**  
**HAZCHEM 2X**  
**In a transport emergency dial 111, Police or Fire Service.**  
**For specialist advice in emergency only, call**  
**0800 243 622 (24 hours).**



**HSNO classification: 6.1C, 6.9A, 8.3A, 9.1B, 9.2A, 9.3B**

### TOXICITY

Toxic – may be fatal if swallowed, inhaled or absorbed through the skin. May cause organ damage from repeated oral exposure at high doses.

### CORROSIVE

Danger – this product is corrosive and may cause eye damage.

### ECOTOXIC

Toxic to aquatic organisms. Avoid contamination of any water supply with product or empty container.

Selective herbicide – very toxic to some plant species  
Beware: Spray drift hazard. Apply this product carefully. Spray drift may cause serious damage to other desirable plants. Ensure compliance with relevant regional air management plans. To minimise the risk of damage from drift of this product, especially where susceptible vegetation is within 0.5km, use low drift nozzles which minimise the production of small droplets (less than 100 microns). Maintain a water rate of at least 30 litres/ha for aerial spraying. Do not add additional wetter unless specified on the label. Preferred wind conditions for spraying are a light constant breeze away from any sensitive crop.

Toxic to terrestrial vertebrates.

### HANDLING PRECAUTIONS

#### STORAGE

Store in original container tightly closed and in a locked, dry, cool area away from foodstuffs, seeds, and fertilisers. Storage must be in accordance with NZS 8409 Management of Agrichemicals.

**Resistant Weeds Warning:**

GROUP	01	HERBICIDE
-------	----	-----------

Resistance to this herbicide could develop from excessive use. To minimise this risk use strictly in accordance with label instructions. Ken-Amine 625 Selective Herbicide contains and belongs to the phenoxy group of chemistry. Ken-Amine 625 Selective Herbicide has the disruption of plant cell growth mode of action. Naturally occurring weed biotypes, resistant to Ken-Amine 625 Selective Herbicide and other phenoxy herbicides, are known to exist. Species with known resistant biotypes include nodding thistle, giant buttercup and slender winged thistle. Resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. Resistant weeds will not be controlled by Ken-Amine 625 Selective Herbicide or other phenoxy herbicides. To prevent or delay the development of resistance use Ken-Amine 625 Selective Herbicide in tank mixtures where appropriate and/or in rotation with herbicides having a different mode of action against the weed.

**GENERAL COMMENTS:**

Ken-Amine 625 Selective Herbicide is damaging to clovers, especially red, suckling and subterranean clover, and plantain and chicory. To minimize clover damage where clovers are actively growing, hard graze to remove as much clover leaf as possible then leave for a few days to freshen up before spraying. Do NOT spray hay or silage paddocks.

After spraying with Ken-Amine 625 Selective Herbicide, poisonous plants (eg Ragwort) may become more palatable. Stock should be kept away from the area until the plants have died down.

Ken-Amine 625 Selective Herbicide will control a wide range of pasture weeds, including thistles and ragwort. Weeds are best controlled as small seedlings. Use the lower rate on seedlings and the higher rate on larger weeds, or during periods of hardened growth. Do not apply if crop or weeds are stressed. Weed control may be unsatisfactory where low soil temperatures inhibit plant growth. Larger plants and established biennials are often not well controlled, and perennial plants are not well controlled. Buttercups and thistles are generally better controlled with MCPA 750. Some weeds (eg Californian thistle, bindweed) require repeat applications.

Rainfall within 6 hours of application may reduce effectiveness.

To improve control of large thistles, or phenoxy resistant thistles add Clopyralid 300g/L at 100ml/ha. Clopyralid 300g/L will severely suppress clover growth and clovers may take 6-12 months to recover.

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residues Limits of Agricultural Compounds) Food Standards.**

Direction for use:

Crop	Rate L/ha	Note
Wheat Barley Millet Ryecorn	1.3-2.0	Apply from 6 leaf stage to jointing (when nodes are visible at the base of the stem). Do not apply to wheat or barley between jointing and flowering. Rapidly growing spring sown cereals should be treated with MCPA 750.  Critical comment: Application to crops outside recommended growth stages or crops stressed due to insect damage or moisture stress may cause serious injury including yield loss. Inspect crops before application to check condition and growth stage.
Oats	1.3	Use only on autumn sown oats and treat in the following spring when plants are fully tillered. Use MCPA 750 on spring sown oats.
Pasture	2-4	Use 2L/ha for seedlings and small plants and 4L/ha for more mature plants. Treat seedling weeds in late autumn/early winter. Graze area to be treated to expose weeds and reduce clover leaf. Then leave for one week to freshen up before spraying. As treated weeds are more palatable to stock, avoid grazing for 7 days after spraying.

**Weeds Controlled in Cereals:**

At 1.3L/ha – fathen, hedge mustard, nettles, shepherd’s purse, wild turnip.

At 2L/ha – black nightshade, calandrinia, docks (seedlings), thistles.

Checked at 2L/ha – chamomiles, cornbind, regrowth docks, fumitory, spurrey, stinking mayweed, willow weed, wireweed, yarrow, vetch. For best results use an alternative cereal herbicide.

**Plant Back Periods:** 2,4-D has soil activity which may be exacerbated by dry soil conditions. To ensure that susceptible species (e.g. brassicas, grasses, clovers) are not damaged by 2,4-D soil residues, do not plant for 4 weeks after application.

**Aerial Application:** Read Spray Drift Hazard section of this label.

**High Volume Spot Treatment:** Use 650ml/100 litres water with a pressure of 600-800kPa and a No. 4 tip and apply as a fine wetting spray. Spray marker dye can be used to identify treated plants.

**Knapsack:** Use 65ml/10 litres water