

HARMFUL • ECOTOXIC

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

KEN-ESTER LV 680

SELECTIVE HERBICIDE

**ACTIVE INGREDIENT: 680 g/L 2,4-D as the 2-ethyl hexyl ester
in the form of an emulsifiable concentrate**

For broadleaf weed control in pasture and turf.



Registered pursuant to ACVM Act 1997, Approval No. P8710
See www.foodsafety.govt.nz for registration conditions
Approved pursuant to HSNO Act 1996, Approval No. HSR000962
See www.epa.govt.nz for approval conditions

Kenso Corporation (M) Sdn Bhd Office A, 49B, Apollo Drive, Rosedale, Auckland 0632 New Zealand Ph: 09 410 0861

KEN-ESTER LV 680 SELECTIVE HERBICIDE

APPROVED HANDLER

This product must be under the control of an approved handler when applied in a wide dispersive manner or used by a commercial contractor.

This product must not be used for any purpose, or in any manner, contrary to this label unless authorised under appropriate legislation.

HSNO CLASSIFICATION: 6.1D, 6.3B, 6.5B, 6.9A, 9.1A, 9.2A, 9.3C

TOXICITY

Harmful - may be harmful if swallowed, inhaled or absorbed through the skin. May cause mild skin irritation. May cause sensitisation from prolonged skin contact. May cause organ damage from repeated oral exposure at high doses.

ECOTOXIC

Very toxic to aquatic organisms with long-lasting effects. Avoid contamination of any water supply with product or empty container. Application to drains, ditches and irrigation channels must only be made when these waterways are dry and at least one month is allowed to elapse

before the channels are used to carry water.

Selective herbicide - very toxic to some plant species.

Beware - Spray drift hazard. Apply this product carefully. Spray drift may cause serious damage to other desirable plants. Ensure compliance with relevant regional air management plans.

To minimise the risk of damage from drift of this product, especially where susceptible vegetation is within 0.5km, use low drift nozzles which minimise the production of small droplets (less than 100 microns). Maintain a water rate of at least 30 litres/ha for aerial spraying. Do not add additional wetter unless specified on the label. Preferred wind conditions for spraying are a light constant breeze away from any sensitive crop. Vapour loss of the product could possibly occur under certain environmental conditions from a sprayed area (soil or vegetation) which has a temperature above 20°C. Avoid using it near sensitive crops on sunny days when the air temperature is forecast above 20°C or on overcast days when the air temperature is forecast to be over 25°C.

FIRST AID

If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If splashed

in eyes, wash out immediately with water. If inhaled, move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped.

SAFETY DATA SHEET

A Safety Data Sheet is available from Kenso Corporation (M) Sdn Bhd on request.

STORAGE

Store in original container tightly closed and in a locked, dry, cool area away from foodstuffs, seeds, and fertilisers. Storage must be in accordance with NZS 8409 Management of Agrichemicals.

Stores containing 100 litres of Ken-Ester LV 680 Selective Herbicide require signage and secondary containment, and more than 100 litres require emergency response plans.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.



EMERGENCY RESPONSE
Ring 0800 CHEMCALL, or
0800 243 622 (24 hours)

NET CONTENTS: 20L



Product of Malaysia



Batch No.:
Date of Manufacture:

UN NO: 3082

ENVIRONMENTALLY HAZARDOUS
SUBSTANCE, LIQUID N.O.S.

(CONTAINS 2,4-D Ester)

PACKING GROUP III

HAZCHEM 2X



In a transport emergency dial 111,
Police or Fire Service.

For specialist advice in emergency only,
call 0800 243 622 (24 hours)

HARMFUL • ECOTOXIC
KEEP OUT OF REACH OF CHILDREN
READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

KEN-ESTER LV 680

SELECTIVE HERBICIDE

**ACTIVE INGREDIENT: 680 g/L 2,4-D as the 2-ethyl
hexyl ester in the form of an emulsifiable concentrate.**

For broadleaf weed control in pasture and turf.

Registered pursuant to ACVM Act 1997, Approval No. P8710
See www.foodsafety.govt.nz for registration conditions

Approved pursuant to HSNO Act 1996, Approval No.
HSR000962
See www.epa.govt.nz for approval conditions



Kenso Corporation (M) Sdn Bhd
Office A, 49B, Apollo Drive
Rosedale, Auckland 0632 NZ
Phone: (09) 410 0861

APPROVED HANDLER

This product must be under the control of an approved handler when applied in a wide dispersive manner or used by a commercial contractor.

This product must not be used for any purpose, or in any manner, contrary to this label unless authorised under appropriate legislation.



HSNO classification: 6.1D, 6.3B, 6.5B, 6.9A, 9.1A, 9.2A, 9.3C
TOXICITY

Harmful - may be harmful if swallowed, inhaled or absorbed through the skin. May cause mild skin irritation. May cause sensitisation from prolonged skin contact. May cause organ damage from repeated oral exposure at high doses.

ECOTOXIC

Very toxic to aquatic organisms with long-lasting effects. Avoid contamination of any water supply with product or empty container. Application to drains, ditches and irrigation channels must only be made when these waterways are dry and at least one month is allowed to elapse before the channels are used to carry water.

Selective herbicide - very toxic to some plant species.

Beware - Spray drift hazard. Apply this product carefully. Spray drift may cause serious damage to other desirable plants. Ensure compliance with relevant regional air management plans.

To minimise the risk of damage from drift of this product, especially where susceptible vegetation is within 0.5km, use low drift nozzles which minimise the production of small droplets (less than 100 microns). Maintain a water rate of at least 30 litres/ha for aerial spraying. Do not add additional wetter unless specified on the label. Preferred wind conditions for spraying are a light constant breeze away from any sensitive crop.

Vapour loss of the product could possibly occur under certain environmental conditions from a sprayed area (soil or vegetation) which has a temperature above 20°C. Avoid using it near sensitive crops on sunny days when the air temperature is forecast above 20°C or on overcast days when the air temperature is forecast to be over 25°C.

PERSONAL PROTECTION

Avoid contact with skin or eyes and inhalation of spray mist. When handling wear eye protection, waterproof gloves, boots and overalls. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work. Wash protective clothing daily after work.

EQUIPMENT

Apply using well maintained and accurately calibrated equipment.

DISPOSAL

Triple rinse empty container and add residue to spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise, crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.

Spillage:

Wear appropriate protective clothing and prevent material from entering waterways. Absorb spill with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely in an approved landfill.

Transport:

Do not carry this product on a passenger service vehicle.

FIRST AID

If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If splashed in eyes, wash out

immediately with water. If inhaled, move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped.

SAFETY DATA SHEET

A Safety Data Sheet is available from Kenso Corporation (M) Sdn Bhd on request.

CONDITION OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

* Not a trademark of Kenso Corporation (M) Sdn Bhd.

Registered pursuant to ACVM Act 1997, Approval No. P8710
See www.foodsafety.govt.nz for registration conditions

Approved pursuant to HSNO Act 1996, Approval No. HSR000962
See www.epa.govt.nz for approval conditions

HANDLING PRECAUTIONS

STORAGE

Store in original container tightly closed and in a locked, dry, cool area away from foodstuffs, seeds, and fertilisers. Storage must be in accordance with NZS 8409 Management of Agrichemicals. Stores containing 100 litres of Ken-Ester LV 680 Selective Herbicide require signage and secondary containment, and more than 100 litres require emergency response plans.

Record Keeping:

Records, as described in NZS 8409 Management of Agrichemicals, must be kept whenever Ken-Ester LV 680 Selective Herbicide is used.

Shelf Life: When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

Mixing:

Pour the measured quantity of Ken-Ester LV 680 Selective Herbicide into the partially filled spray tank and with the agitator running top up with the remainder of the water.

Cleaning:

Flush equipment thoroughly with clean water after use. Decontaminate before using on plants that are sensitive to 2,4-D.

Decontamination:

After thoroughly washing the tank and flushing the system with clean water, quarter fill the tank, add 500g of an alkali detergent and circulate throughout the system for 15 minutes. Drain the system. Remove the filters and nozzles and clean them separately. Then flush the system with clean water and allow to drain.

Compatibility: Ken-Ester LV 680 Selective Herbicide is compatible with many other agricultural chemicals including Dicamba 500g/L and Clopyralid 300g/L.

Resistant Weeds Warning:

Naturally occurring biotypes resistant to Ken-Ester LV 680 Selective Herbicide and other phenoxy herbicides are known to exist. These become more prevalent after many years of continuous use of the product. Species with known resistant biotypes include nodding thistle, giant buttercup and slender winged thistle. To prevent or delay the development of resistant weeds use Ken-Ester LV 680 Selective Herbicide in tank mixes where appropriate, and/or in rotations with herbicides having a different mode of action against the weed.

Batch No. :

Date of Manufacture :

EMERGENCY RESPONSE
Ring 0800 CHEMCALL, or
0800 243 622 (24 hours)



UN NO : 3082
ENVIRONMENTALLY HAZARDOUS SUSTANCE,
LIQUID N.O.S.
(CONTAINS 2,4-D Ester)
PACKAGING GROUP III
HAZCHEM 2X
In a transport emergency dial 111, Police or Fire Service.
For specialist advice in emergency only, call
0800 243 622 (24 hours).

GENERAL COMMENTS:

Ken-Ester LV 680 Selective Herbicide is most effective when applied to actively growing seedling weeds. Established weeds, particularly perennial species, may not be well controlled. Do not apply to plants under stress from frost or drought. Buttercups and thistles are generally better controlled with MCPA 750. Some weeds (e.g. Californian thistle, bindweeds) require repeat applications.

Application Timing: Autumn, early winter or early spring application is recommended for the best control of weeds. Cold temperatures and frosts limit plant growth and resultant weed control from applications during mid winter months may be unsatisfactory.

Pasture Management: Ken-Ester LV 680 Selective Herbicide damages plantain, chicory and clover, especially red, subterranean and suckling clover. Damage is temporary and recovery of perennial clovers usually occurs after several months. To minimise clover damage graze pasture before spraying to remove clover leaf and expose weeds. Do not graze so hard that weed foliage is removed. Let pasture freshen for one week before spraying. Do not graze for two weeks after application to allow maximum uptake and translocation of the herbicide. Premature grazing may reduce weed control. Light rainfall 2 hours after application should not adversely affect results. After spraying with Ken-Ester LV 680 Selective Herbicide, poisonous plants (e.g. ragwort, hemlock) may become more palatable and stock should be kept away until the plants have died down.

Plant Back Period: 2,4-D has soil activity which may be exacerbated by dry soil conditions. To ensure that susceptible species (e.g. brassicas, grasses or clovers) are not damaged by 2,4-D soil residues, do not plant for 4 weeks after application.

Application: Ken-Ester LV 680 Selective Herbicide can be applied using conventional ground or aircraft spraying equipment. Boom Spraying: Use 100-300 litres of water/ha. For difficult to control perennial pasture weeds use 300-500 litres of water/ha to improve coverage. CDA Spraying - Pasture Weeds: Use 15-50 litres of water/ha. Droplet size must exceed 150 microns. Aerial Spraying: For pasture weeds apply with calibrated equipment in 30-60 litres of water/ha. Spot Spraying: For pasture weeds use a pressure of 600-800kPa through a No. 4 tip and apply as a fine misting spray. Complete coverage is essential. Use 500ml/100 litres of water. For a knapsack use 5ml/litre.

Weeds Controlled:

Ken-Ester LV 680 Selective Herbicide controls or severely checks the following: Alligator weed, allseed, apple of Peru, Australian flax, Australian sedge, bachelor's button, Bathurst bur, beggar's ticks, bitter cress, black nightshade, burdock, bur medick, buttercups, Californian poppy, Canadian fleabane, Cape tulip, castor oil plant, catsear, charlock, chicory, clovers, cornbind, cresses, cudweed, dandelion, dove's foot, fathen, field bindweed, field cress, foxglove, goat's rue, greater bindweed, groundsel, hawkbit, hawksbeard, hawkweed, hedge mustard, hemp nettle, hoary cress, horehound, marsh cress, Mexican devil, mint weed, monkey musk, mouse-ear chickweed, nettles, oxtongue, parsley dropwort, penny cress, piripiri, plantains, poplars, poroporo, ragwort, red dead-nettle, redroot, rushes (leafless), selfheal, shepherd's purse, silver poplar, Spanish heath, storksbill, tansy, teasel, thistles (nodding, scotch, variegated, welter, winged), twin cress, vetch, wart cress, wild carrot, wild mustard, wild parsnip, wild radish, wild turnip, willows, willow weed, wineberry.

Weeds controlled as seedlings only: Cape weed, creeping mallow, cut-leaved geranium, daisies, docks, fennel, fumitory, inkweed, lupin, Onehunga weed, wild portulaca, spurrey, storksbill, tutu, viper's bugloss.

Ken-Ester LV 680 Selective Herbicide does not control: Barberry, bracken, chamomile, chickweed, cleavers, grasses, hawthorn, hydrocotyle, stinking mayweed, oxalis, prostrate amaranth, parsley piert, pearlwort, sorrel, staggerweed, stonecrop, tutsan, and many semi-woody species after the seedling stage.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residues Limits of Agricultural Compounds) Food Standards.

Directions For Use:

Established pastures and waste areas: Apply 750ml – 3L/ha. Refer to the weeds controlled tables for guidance.

Thistles			
	Rate L/ha	Stage	Notes
Nodding thistle	1.5	Seedlings	Root crowns less than 2cm diameter
	2.0-2.3	Rosettes	Root crown 2-4cm diameter, before flower stalk develops
Scotch thistle	0.75	Seedlings	
	1.5	Rosettes	
Variegated thistle	1.5	Seedlings	
	2.0-2.3	Rosettes	Less than 15cm diameter
Winged thistle	1.5	Seedlings	
	2.0-2.3	Rosettes	Prior to flower stalk development

To improve control of large thistles, or phenoxy resistant thistles add Clopyralid 300g/L at 100ml/ha. Clopyralid 300g/L will severely suppress clover growth and clovers may take 6-12 months to recover.

Other Broadleaf Weeds			
	Rate L/ha	Stage	Notes
Annual buttercups, bur medick, catsear, cresses, dandelion, hedge mustard, hemlock, plantains, redroot	0.75	Seedlings	
	1.5	Small rosettes	
Charlock, wild radish, shepherd's purse, wild turnip	1.5	Seedlings	
	2.3-3.0	Small rosettes	
Apple of Peru, Bathurst bur, black nightshade, burdock, creeping mallow, docks, field bindweed, fireweed, fennel, foxglove, hoary cress, nettles, parsley dropwort, storksbill, vetch, wild portulaca, wild carrot, willow weed	2.0-3.0	Seedlings only	Docks: regrowth from root chips is not controlled. Vetch: spray at multiple leafstage.
Ragwort	2.0	Seedlings	Large rosettes or multicrowns require spot treatment with other herbicides.
	3.0	Small rosettes	

Woody and Non-Crop/Pasture Weeds		
	Rates	Notes
Willows	Aerial Spraying: 6L in 200-400L water/ha	Apply in late spring to fully leafed trees. A second treatment may be required 12 months later.
	Gun Spraying: 350ml/100L water	Apply in late spring to fully leafed trees. Ensure complete coverage.
	Stump swabbing or frilling: Use a 1:40 mixture of Ken-Ester LV 680 Selective Herbicide in diesel or waste oil.	Apply to cut or frilled surface immediately after cutting using a paint brush or knapsack.
Spanish heath	500ml/100L water	Apply to complete coverage during active spring growth.
Cape tulip, goat's rue seedlings, and leafless rushes	3-6L/ha in 300-500L water as an overall boom spray	Treat rushes with the high rate in spring.
Australian sedge	530ml plus 100ml paraquat (25%) / 100L water	Spot spray in the April-November period. Ensure plants are completely wet.