

**KEEP OUT OF REACH OF CHILDREN
READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING**

SULPHUR 800 WG

FUNGICIDE/MITICIDE

ACTIVE INGREDIENT: 800 g/kg SULPHUR in the form of a water dispersible granule.

For the control of certain diseases of Apples, Grapes, Stonefruit, Tamarillos,
Vegetables and Ornamentals.



Registered pursuant to ACVM Act 1997, Approval No. P8992
See www.foodsafety.govt.nz for registration conditions
Approved pursuant to HSNO Act 1996, Approval No. HSR000741
See www.epa.govt.nz for approval conditions

NET CONTENTS: 15KG

Registrant:
Kenso Corporation (M) Sdn Bhd
Lot 29, Jalan Sungai Pinang 5/4, Taman Perindustrian
Pulau Indah, 42920 Pelabuhan Klang, Selangor, Malaysia.

NZ Agent:
Kenso NZ Limited
2 Bond Crescent, Forrest Hill, Auckland 0620 New Zealand.
Ph: 0800 537 766

HSNO CLASSIFICATION: 6.4A, 9.1D**WARNING**

Harmful - May cause eye irritation. Harmful to aquatic organisms.

PRECAUTIONS**STORAGE**

Keep out of reach of children. Store in the original packaging away from foods. Storage of certain quantities of this product trigger specific storage requirements (Refer to HazNote). When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.

PERSONAL PROTECTION

Avoid contact with eyes and skin. Avoid inhalation of dust and spray mist. When mixing or applying, wear overalls, chemical resistant boots and gloves, head covering and eye protection. Clean protective clothing daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

EQUIPMENT

Apply with well-maintained and calibrated spray equipment.

DISPOSAL

Dispose of this product only by using in accordance with this label, or at a suitable landfill. Do not burn. Dispose of packaging to a suitable landfill. Do not burn. Do not use packaging for any other purpose.

ENVIRONMENTAL

Use this product carefully. Do not contaminate aquatic environments with product, spray drift or packaging. Do not allow spray drift outside the target area to occur.

SPILLAGE

Exclude non-essential people from the area. Wear protective clothing. Moisten to prevent dusting and sweep up. Place in a waste container. Dispose of safely to an approved landfill.

GENERAL INFORMATION

Sulphur 800 WG Fungicide/Miticide is a protectant fungicide and must be applied before disease infection occurs. For best results, even spray coverage is necessary.

CAUTION

Some fruits and crops are susceptible to injury from sulphur under certain climatic conditions. The user is advised not to use on any crop unless local experience has proven sulphur to be safe in that locality. Do NOT apply Sulphur 800 WG Fungicide/Miticide when the temperature is above 25°C, or under conditions of extreme humidity, as crop damage can occur. Cucurbits and other crops grown under glass can be affected by an application of Sulphur 800 WG Fungicide/Miticide during periods of rapid growth. Do NOT apply to sulphur shy apples such as Cox's Orange.

Do NOT apply to apricots.

Do NOT apply to pears.

Do NOT apply to apples with captan or within 5 days of a captan treatment.

Do NOT apply with crop oil or within 21 days of a crop oil application.

APPLICATION METHODS

Suitable for use with all applicators. The rates expressed are for high volume spraying to run off. For concentrated spraying, adjust dilution rate accordingly. Sulphur is corrosive so all spray equipment should be thoroughly flushed with clean water after use.

COMPATIBILITY

Sulphur 800 WG Fungicide/Miticide is compatible with most insecticides, fungicides and foliar fertilisers. When using with untried emulsifiable concentrate formulations, test the compatibility of both products by mixing small quantities before tank mixing. Do NOT use with spraying oils or any products containing mineral oil.

RAINFALL

Rain within 3 hours of treatment may reduce efficacy.

WITHHOLDING PERIODS

Nil.

FIRST AID

If swallowed do not induce vomiting. If splashed in eyes, wash out immediately with running water for several minutes. If skin or hair contact occurs, remove contaminated clothing and wash with soap and running water. If inhaled, immediately move to fresh air and rest. Maintain half upright position if breathing is difficult. Call the National Poisons Centre 0800 POISON (0800 764 766) (24 hrs) or a doctor immediately for advice.

SAFETY DATA SHEET

A Safety Data Sheet is available from the Kenso NZ website: www.kenso.co.nz

It is an offence to use to use this product on animals.

DIRECTIONS FOR USE

Crop	Pest /Disease	Rate	Remarks
Apples	Powdery mildew	100-200 g per 100 litres	Apply from bud movement until harvest, however, use before December may affect fruit finish. DO NOT apply in temperatures above 25°C. DO NOT apply in combination with captan. DO NOT apply to sulphur sensitive varieties such as Kidds Orange Red, Cox's Orange Pippin or Red Delicious. Fruit finish may be affected on some apple varieties. DO NOT apply on pears.
Grapes	Powdery mildew	150-300 g per 100 litres	Apply at 10 to 14 day intervals from bud swell until 6 weeks before harvest. Use the higher rates when disease is evident. Erineum mites will also be controlled.
Peas	Powdery mildew	100-200 g per 100 litres	Apply when conditions favour disease and repeat as necessary. Use the higher rates where infection has already occurred.
Peaches Plums Nectarines	Rust	100-200 g per 100 litres	Apply at fruit set then at 21 day intervals until harvest. Use the higher rate during conditions of high pressure. DO NOT apply to apricot.
Roses	Powdery mildew	200-300 g per 100 litres	Apply on a 14 day schedule using the higher rate during periods of severe disease pressure. DO NOT use in mixture with maldison.
Tamarillos	Powdery mildew	100-200 g per 100 litres	Apply at 10 to 14 day intervals throughout the growing season.
Vegetables	Powdery mildew	100-300 g per 100 litres or 2-3kg/ha	Use at regular 10 - 14 day intervals during periods of infection. Use the higher rate where infection has already occurred.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.



EMERGENCY RESPONSE
Ring 0800 CHEMCALL, or
0800 243 622 (24 hours)

Product of India



Batch No.:

Date of Manufacture: