

DANGER

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

KEN-UP DRY 680 SG HERBICIDE

ACTIVE INGREDIENT: 680 g/kg GLYPHOSATE
as the mono-ammonium salt in the form of a water-soluble granule

GROUP **9** HERBICIDE

A highly concentrated granular Glyphosate suitable for the control of a wide range of broadleaf weeds and grasses in agriculture, horticulture, forestry and non-cropland areas.

Registered pursuant to the ACVM Act 1997, No. P008339
See www.foodsafety.govt.nz for registration conditions
Approved pursuant to the HSNO Act 1996, Approval No. HSR000219
See www.epa.govt.nz for approval controls



NET CONTENTS: 10 KG

Registrant:
Kenso Corporation (M) Sdn Bhd
Lot 29, Jalan Sungai Pinang 5/4, Taman Perindustrian Pulau
Indah, 42920 Pelabuhan Klang, Selangor, Malaysia.

NZ Agent:
Kenso NZ Limited
2 Bond Crescent, Forrest Hill, Auckland 0620 New Zealand.
Ph: 0800 536 766

KEN-UP DRY 680 SG HERBICIDE

PERSONNEL QUALIFICATION & EXPERIENCE

This product must be under the control of an appropriately trained and experienced person during storage, use and disposal. For guidance refer to the requirements for class 6 and 9 substances of the Health and Safety at Work (Hazardous Substances) Regulations 2017 part 4.5 and the Hazardous Substances (Hazardous Property Controls) Notice 2017 Part 4 Subpart C.

HAZARD CLASSIFICATION

Serious eye damage Category 1, Hazardous to the aquatic environment chronic Category 2.

DANGER

This product is corrosive and may cause eye damage.

Toxic to aquatic organism with long lasting effects. Avoid contamination of any water supply with product or empty container.

Non-selective Herbicide - toxic to plants. Avoid contact with foliage or green stems of fruit or crops, desirable plants and trees, since severe injury or death may occur.

BEWARE: Apply this product carefully. Spray drift may cause serious damage to other desirable plants.

FIRST AID

If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If splashed in eyes, wash out immediately with water. If inhaled, move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped.

SAFETY DATA SHEET

A Safety Data Sheet is available from the Kenso NZ website: www.kenso.co.nz

STORAGE

Keep out of reach of children. Store under lock and key, in original container, tightly closed, away from human and animal foodstuffs, medicines and remedies, insecticides, fungicides seeds and fertilisers. Segregate from incompatible hazardous substances (Classes 1, 4 & 5). Store in a cool, dry, well ventilated place and protect from sunlight.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.



EMERGENCY RESPONSE
Ring 0800 CHEMCALL, or
0800 243 622 (24 hours)

Product of China



9 421028 660914



Batch No.:
Date of Manufacture:

UN NO: 3077
ENVIRONMENTALLY HAZARDOUS
SUBSTANCE, SOLID, N.O.S.
(CONTAINS GLYPHOSATE)
PACKING GROUP III, HAZCHEM 2X

In a transport emergency dial 111, Police or
Fire Service. For specialist advice in emergency
only, call 0800 243 622 (24 hours)



DANGER

KEEP OUT OF REACH OF CHILDREN
READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

KEN-UP DRY 680 SG HERBICIDE

ACTIVE INGREDIENT: 680 g/kg GLYPHOSATE as the mono-ammonium salt in the form of a water-soluble granule.

GROUP **9** HERBICIDE

A highly concentrated granular Glyphosate suitable for the control of a wide range of broadleaf weeds and grasses in agriculture, horticulture, forestry and non-cropland areas.

Registered pursuant to the ACVM Act 1997, No. P008339
See www.foodsafety.govt.nz for registration conditions
Approved pursuant to the HSNO Act 1996, Approval No. HSR000219
See www.epa.govt.nz for approval controls



Kenso Corporation (M) Sdn Bhd
2 Bond Crescent , Forrest Hill
Auckland 0620 NZ
Phone: 0800 536 766

PERSONNEL QUALIFICATION & EXPERIENCE

This product must be under the control of an appropriately trained and experienced person during storage, use and disposal. For guidance refer to the requirements for class 6 and 9 substances of the Health and Safety at Work (Hazardous Substances) Regulations 2017 part 4.5 and the Hazardous Substances (Hazardous Property Controls) Notice 2017 Part 4 Subpart C.



HAZARD CLASSIFICATION

Serious eye damage Category 1, Hazardous to the aquatic environment chronic Category 2.

DANGER

This product is corrosive and may cause eye damage.
Toxic to aquatic organism with long lasting effects. Avoid contamination of any water supply with product or empty container.
Non-selective Herbicide - toxic to plants. Avoid contact with foliage or green stems of fruit or crops, desirable plants and trees, since severe injury or death may occur.

BEWARE: Apply this product carefully. Spray drift may cause serious damage to other desirable plants.

MIXING INSTRUCTIONS

Ken-Up Dry 680 mixes readily with water. Reduced results may occur if dirty water is used (**e.g. water from ponds or unlined ditches**) or if hard water containing calcium salts is used. Ensure the spray tank is free of any residue of previous spray materials.

Procedure

Premix Ken-Up Dry 680 with water into a slurry prior to adding to a half full spray tank. If adding other herbicides add prior to adding Ken-Up Dry 680 to the tank. Use tank mixtures promptly. If using Silken add last to avoid foaming.

GENERAL INFORMATION

Ken-Up Dry 680 is a broad spectrum, non-selective, non-residual herbicide that is non-volatile and may be used in agriculture, horticulture and for other uses including general weed control. Ken-Up Dry 680 is recommended prior to sowing pasture and crops. Ken-Up Dry 680 is absorbed by plant foliage and green stems. It is inactivated immediately upon contact with the soil and does not provide residual weed control. Ken-Up Dry 680 moves from the point of foliage or stem contact into the root system controlling the plant above and below the ground. Visible effects may take up to 14 days or even longer depending on growing and weather conditions. These effects are a gradual wilting and yellowing of the foliage which advances to deterioration of above and below ground parts. Avoid contact with foliage or green stems of desirable plants and trees since severe injury or destruction may result. Avoid application in winds which allow drift onto desirable plants. Apply Ken-Up Dry 680 only to healthy, actively growing plants with adequate leaves for herbicide uptake. Avoid treating plants that are drought affected, frost damaged, silt laden, or suffering the effects of previous herbicide application. Rainfall occurring up to 6 hours after application may reduce effectiveness. When Silken is added the rainfree period may be reduced to 30 minutes. If Ken-Up Dry 680 is applied to wet plants, the period between application and rainfall, without affecting performance, will be longer. After application of Ken-Up Dry 680 a withholding period for grazing stock is not required (except where Ragwort is present). However, it is recommended that grazing is delayed for one day after treatment of annual weeds or three days if perennial weeds are present to ensure absorption of Ken-Up Dry 680.

USE RECOMMENDATIONS

Refer to "Weeds Controlled" section for rates of application.

General Weed Control:

Handgun - mix 0.5 kg of Ken-Up Dry 680 with 100 litres of water.
Knapsack - mix 75 g of Ken-Up Dry 680 with 15 litres of water.
Apply to completely wet the foliage but not to run-off.

Brushweed Control:

Ken-Up Dry 680 may be used to control a wide range of brushweeds and woody plants.

Agricultural Areas:

Ken-Up Dry 680 may be used for the control of annual and perennial weeds as directed in agricultural land prior to sowing any edible or non-edible crop or pasture. Where White Clover control is required, tank mix with a suitable White Clover herbicide and observe recommended waiting periods prior to planting legumes. Cultivation may start from 7 days after spraying perennial weeds and 3 days for annual weeds.

Pasture: Direct (Spot) Application

Ken-Up Dry 680 is non-selective and may damage or kill any plant in the sprayed area. Retreatment and/or pasture improvement may be necessary to restrict seedling re-establishment.

Selective Wiper Application:

For the control of erect weeds in pastures.

STORAGE

Keep out of reach of children. Store under lock and key, in original container, tightly closed, away from human and animal foodstuffs, medicines and remedies, insecticides, fungicides seeds and fertilisers. Segregate from incompatible hazardous substances (Classes 1, 4 & 5). Store in a cool, dry, well ventilated place and protect from sunlight.

FIRST AID

If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If splashed in eyes, wash out immediately with water. If inhaled, move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped.

SAFETY DATA SHEET

A Safety Data Sheet is available from Kenso NZ website www.kenso.co.nz

CONDITION OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

Registered pursuant to the ACVM Act 1997, No. P008339
See www.foodsafety.govt.nz for registration conditions
Approved pursuant to the HSNO Act 1996, Approval No. HSR000219
See www.epa.govt.nz for approval controls

Batch No. :

Date of Manufacture :

EMERGENCY RESPONSE
Ring 0800 CHEMCALL, or
0800 243 622 (24 hours)

UN NO : 3077



**ENVIRONMENTALLY HAZARDOUS
SUSTANCE, SOLID, N.O.S.
(CONTAINS GLYPHOSATE)
PACKAGING GROUP III
HAZCHEM 2X**

In a transport emergency dial 111, Police or Fire Service.

**For specialist advice in emergency only, call
0800 243 622 (24 hours).**

Pasture Renovation:

Ken-Up Dry 680 may be used to control existing pasture species prior to re-seeding pasture or establishing other crops.

Pre-Harvest: Wheat, Barley, Oats, Peas

For Wheat, Barley and Oats - apply 7–12 days prior to harvest when the grain moisture content is less than 30%.
Peas - Apply 7–14 days prior to harvest when the pods have dried. Use only on peas for threshing. Do not harvest within 3 days after treatment to allow time for the herbicide to translocate into the weed roots.

Do not apply to undersown crops including Clovers as they may be severely damaged.

Do not apply to crops which will be harvested for seed.

Use rates up to 2 kg/ha depending on the specific weeds to be treated. Avoid treating drought stressed plants.

Orchards/Vines: Citrus, Apple, Pear, Stonefruit, Kiwifruit, Grapevines.

Apply as a directed or shielded spray or with wiper equipment. For long term control of annual weeds Ken-Up Dry 680 may be tank mixed with a suitable residual herbicide. Do not apply near unprotected trees less than 3 years old. Do not allow spray or spray drift to contact any part of stone fruit trees, or green bark, suckers, buds, foliage or fruit of other trees. Do not allow wiper surfaces to contact any part of the trees. Weed foliage coated in copper or calcium based compounds may reduce the activity of Ken-Up Dry 680.

Market Gardens:

Ken-Up Dry 680 can be applied before or after seeding and prior to crop emergence. Ken-Up Dry 680 can be applied prior to transplanting or as a post harvest clean-up. For longer term control of seedling weeds, Ken-Up Dry 680 may be tank mixed with suitable residual herbicides. Use in accordance with the label recommendations for these products.

Asparagus: Ken-Up Dry 680 can be applied:

– Following the final harvest and immediately after removal of all spears but before crop regrowth occurs.

– Prior to planting or crop emergence. Do not treat emerging spears.

In established ferns use as a directed or shielded spray or with wiper equipment.

Do not allow spray, spray drift or the wiper surface to contact any part of the Asparagus plants.

Industrial and Non-Agricultural Areas:

Ken-Up Dry 680 can be used for the control of annual and perennial weeds as directed.

For residual control Ken-Up Dry 680 can be tank mixed with simazine or another suitable residual herbicide.

Exercise caution when using residual herbicides prior to planting trees, shrubs or other ornamental plants.

Turfgrass and Recreational Areas:

Apply before establishing lawns, parks, reserves and ornamentals. Ken-Up Dry 680 can also be applied around the base of established trees as a directed or shielded spray or by wiper equipment. Do not allow spray or spray drift or the wiper surface to contact any part of the tree. In mown areas, allow sufficient re-growth before spraying.

Aquatic Areas Drains and Waterways

Ensure that 75% of the weed to be treated is above water level or efficacy may be reduced. Following spray application do not submerge the treated plant as this will cause the herbicide to be washed off. Avoid treating silt laden plants.

Forestry:

Ken-Up Dry 680 can be applied for the control of grasses and a wide range of brushweeds.

The addition of Silken is required for the control of bracken and gorse.

Do not allow spray, spray drift or the wiper surface to contact the foliage or green bark of desirable trees as severe injury can occur.

APPLICATION INSTRUCTIONS

Ensure the equipment is in good operating order and correctly calibrated.

Wash thoroughly with water before use, and ensure filters and nozzles are free of pesticide residue.

Avoid spray drift. Care must be used when applying this product to prevent injuring desirable plants and crops.

To clean equipment after use, flush thoroughly with water.

Boom Application:

Water volumes up to 200 litres/ha of clean water are desirable. For broadacre boom application flat fan, wide angle nozzles (e.g. 110°) using pressures of 200–280 kPa are preferred. Set the boom height to ensure a double overlap of the spray pattern at the top of the weed canopy.

For orchard application wide angle, even spray nozzles are preferred.

Knapsack and Handgun Application:

Adjust the equipment to achieve a fine spray. Use the minimum spray volume necessary to achieve uniform and complete coverage without runoff.

Handgun spraying pressures of 1000–2000 kPa and No. 4 to No. 6 spray tips are recommended.

WEEDS CONTROLLED					
WEED	Boom or aerial kg/ha	Handgun kg/100 litres	Knapsack g/15 litres	Mistblower g/10 litres	Critical Comments
Annual Weeds (General)	0.5-1.0	0.1	15	30	Use higher rate for larger weeds.
Australian Sedge	3 Aerial 4.5	0.5	75	150	Apply from November to March. Where rank growth is present the area should be burnt 12–18 months prior to aerial application to ensure adequate spray coverage.
Barbery	-	0.5	75	150	Apply January to April from flowering to late fruit. Where bushes have been cut, regrowth should be 1.5 metres tall before spraying.
Barley Grass	0.5	0.1	15	30	
Blackberry	4.5-10	0.5-0.75	75	150	Apply to healthy plants from blossom to ripe fruit stage prior to Autumn frosts (December to April). Only bushes with a well developed canopy of leaves should be treated. Early treatment (December to January) may be preferred in drought prone areas. Leave sprayed blackberry for 40–60 days before burning or slashing and oversow the area. Where treatment has been carried out under marginal conditions some regrowth can be expected within 1–2 years. For aerial application of hard to kill Blackberry or where no follow up pasture management is anticipated, use the higher rate.
Boxthorn	-	0.7	-	-	Apply November to April to actively growing plants. Avoid salt damaged plants. Spray to ensure complete coverage of dense thickets. Do not use additional surfactant as poor results may occur.
Bracken	4.5 + Silken 250ml/100 litres	0.5 + Silken 200ml/100 litres	75 + Silken 30ml	-	Apply when fronds are unfurled (mid January to June) but prior to frost browning. Dense virgin bracken with high trash content and low leaf to root ratio should be burnt in spring and the unfurled frond regrowth sprayed the following March to June.
Broom	4.5 + Silken 250ml/100 litres	0.5 + Silken 200ml/100 litres	75 + Silken 30ml	150 + Silken 20ml	Aerial Application - Best results are obtained when plants are in full leaf. Handgun Application - Apply all year round. Ensure full coverage.
Browntop	1.5	0.25	40	75	Apply to 5–10 cm high actively growing plants.
Buddleja	4.5	0.5	75	-	Handgun: If actively growing treat all year. Aerial: Treat Buddleja up to 2 metres high during Feb. to Apr.
Californian thistle	2	0.5	75	150	Apply after flower bud development but before autumn browning. To ensure maximum shoot emergence do not cultivate for 6 months prior to treatment. Wiper: A double pass at low speed is recommended in heavy infestations. Mowing thistles 10 days after treatment may assist control. Retreatment may be necessary in the following season.
Cleavers	0.75	0.25	40	75	
Couch	1.5-3	0.5	75	150	For suppression use 1.5 kg/ha, for control use 2–3 kg/ha when growing conditions are satisfactory. Use the high rate in late spring. Apply to plants with at least 4–5 leaves and active shoot and root growth. Do not cultivate for 6 months prior to application to ensure maximum shoot emergence. See the Use Recommendations section for control of Couch in a pre harvest situation. In areas of dense infestation, a repeat application may be necessary.
Dock	2-4.5	0.5	75	-	Treat plants in spring or summer when actively growing and in full leaf. Use low rate for seedlings and young plants. Use high rate for control of established plants.
Floating Sweet Grass/ Reed Sweet Grass	3	0.5	75	150	Apply to actively growing plants when most have reached early seedhead.
Gorse	-	0.5 + Silken	-	-	Apply all year round.

Wiper Application:

Pre-mix 1 part of Ken-Up Dry 680 and 4 parts of clean water. Mix only enough herbicide for immediate needs and allow the wiping area to become fully saturated before commencing. The wiping area must remain moist and clean otherwise reduced results will occur. In heavy weed infestations or when treating perennial weeds such as Californian Thistle or multi-crown Ragwort, a double pass in opposite directions on the same day will give the best results. As weed density increases, decrease ground speed. Drain thoroughly and wash immediately after use.

Aircraft Application:

Ken-Up Dry 680 may be applied by aircraft in forestry and non-cropping areas only. Aerial application of Ken-Up Dry 680 is not recommended on cropping land or near desirable plant species as even slight drift may cause severe injury or destruction. Do not apply in gusty or windy conditions, or in other conditions conducive to drift e.g. still air and inversion conditions.

Mistblower Application:

Use a solution of 150 g of Ken-Up Dry 680 in 10 litres of water. For the micron attachment increase to 1 kg of Ken-Up Dry 680 in 10 litres of water. Provide complete coverage and spray to wet but not to run-off.

SILKEN

Silken may be used with Ken-Up Dry 680 to improve rate of uptake and shorten the rainfree period on certain weeds. Refer to Silken label before using.

WEEDS CONTROLLED					
WEED	Boom or aerial kg/ha	Handgun kg/100 litres	Knapsack g/15 litres	Mistblower g/10 litres	Critical Comments
Indian Doab	4.5	0.5	75	-	Apply when most plants have reached early seedhead.
Johnson Grass	3	0.5	75	150	Apply to actively growing plants.
Kikuyu	3	0.5	75	150	Apply to actively growing plants. Under dense growth conditions silorate and allow regrowth before treatment to improve control levels.
Mercer Grass Aquatic Non-Aquatic	4.5 3	0.75 0.5	100 75	225 150	Apply to actively growing plants. Under dense growth conditions silorate and allow regrowth before treatment to improve control levels.
Nodding Thistle	-	0.5	75	-	Wiper Application (see Application Instructions). Apply when actively growing when at least 10 cm of the weed can be wiped. In heavy infestations a double pass is recommended.
Old Man's Beard	-	0.5-1	150	-	Apply November to March prior to leaf senescence. Vines too high for complete coverage should be cut at ground level in winter or spring and the regrowth sprayed in March when plants are at least 0.5 metres long. Follow-up treatment will be necessary to control seedlings. Use the low rate for small vines.
Pampas Grass/Toetoe	-	0.5	75	-	Apply Spring-Autumn. Ensure complete coverage. Best results are achieved when applied after flowering.
Paspalum	2-3	0.5	75	150	Use the low rate from January to April (early seedhead) prior to frosts. Use 3 kg/ha on vegetative plants.
Potato (self set)	1.5	0.25	40	75	
Prairie Grass	0.75	-	-	-	
Ragwort	3	0.5	75	-	Apply when actively growing. Wiper: Apply when plants are 10–15 cm above the pasture. In heavy infestations a double pass is recommended.
Ragwort Caution: Ragwort plants may become more palatable following spraying and stock should be removed until the treated plants brownout.					
Ratstail	2.5	-	-	-	Apply to actively growing plants in Autumn.
Rautiahi (cutty grass)	3	0.5	75	150	Apply to actively growing plants from January to March. Burning or slashing may take place from 30 days after treatment following brownout.
Ring Fern	0.5-1.5	0.15	25	-	Control: Apply 1.5 kg/ha in January to April. Pasture Manipulation: Apply 0.5 – 1 kg/ha December to April. Use the high rate for dense infestations. Burn 4–6 weeks after treatment and oversow with pasture.
Rushes	3	0.5	75	150	Apply to actively growing plants from January to March. Burning or slashing may take place from 30 days after treatment following brownout. Wiper: Apply November to May.
Ryegrass Perennial	0.75-1.5 + Silken 100ml/100 litres	0.25 + Silken 100ml/100 litres	40 + Silken 15ml	-	Apply any time plants are actively growing and 3–10 cm high. Use low rate January to July when cultivation is to follow spraying Use 1.5 kg/ha August to December.
Sweet Briar	6 + Silken 250ml/100 litres	0.5 + Silken 200ml/100 litres	75 + Silken 30ml	-	Apply from late flowering to early leaf yellowing. Best results achieved with full coverage.
Tall Fescue	3	0.5	75	150	Apply to actively growing plants January to March.
Willow	4.5	0.5	-	-	Apply from January to April before leaf fall begins. Total spray coverage is essential for best results.
Yarrow	1.5-3	-	-	-	Apply to plants with 5–10 cm of leaf. Cultivate with a minimum of 2 grubblings from 10 days after

PRE TREATMENT CHECKLIST

- Do not apply in conditions where drift may occur e.g. wind.
- Weeds should be clean, actively growing and not under stress from drought, frost, cold, water logging, grazing or previous herbicide applications.
- Do not apply Ken-Up Dry 680 if rainfall is likely within 6 hours (Silken may be used to reduce this period).
- Ensure sufficient leaf is present for uptake of Ken-Up Dry 680.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residues Levels for Agricultural Compounds.

POST TREATMENT MANAGEMENT

- A withholding period for grazing stock is not required (except where ragwort is present). It is, however, recommended that grazing or cultivating treated plants is delayed for one day for annual weeds and three days for perennial weeds to ensure absorption and translocation of Ken-Up Dry 680.

SHELF LIFE: When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

RESISTANCE MANAGEMENT **GROUP 9 HERBICIDE**

Ken-Up Dry 680 contains the GROUP 9 herbicide, glyphosate. Do not use another product containing a herbicide in the same group in conjunction with Ken-Up Dry 680, and where possible alternate the types of herbicides used within any one paddock over time. Herbicides from different group codes should either be used alternately or mixed together to prevent the numbers of resistant individuals building up. Refer to <https://resistance.nzpps.org/> for further resistance management advice.